



Ansell**CARES**SM
Education. Evidence. Engagement.

DOUBLE GLOVING

Evidence for best practice

CLINICAL BULLETIN

DOUBLE GLOVING




Double gloving is advocated by many international health and perioperative organizations as listed below. These peak bodies recognize the need to advocate for the benefits and positive outcomes of double gloving. Understanding the evidence-based rationale for double gloving in perioperative practice is key to educating others and improving compliance outcomes.

- AAOS** - American Academy of Orthopedic Surgeons
- ACORN** - Australian College of Perioperative Nurses
- ACS** - American College of Surgeons
- AfPP** - The Association for Perioperative Practice
- AORN** - Association of periOperative Registered Nurses
- AST** - Association of Surgical Technologists
- EORNA** - European Operating Room Nurses Association
- CDC** - Centers for Disease Control and Prevention
- NICE** - The National Institute for Health and Care Excellence
- ORNAC** - Operating Room Nurses Association of Canada
- WHO** - World Health Organization


OVERVIEW

Evidence has shown that surgical glove microperforations occur commonly during surgical procedures regardless of the specialty, although the risk increases with the number of instruments used, the time of wear, and the surgical experience of the wearer. Double gloving has shown to provide overall better protection from microperforations and sharps injury.¹

FACTORS THAT MAY RESULT IN MICROPERFORATIONS



The operating theatre is a fast paced and high stress environment, containing the greatest concentration of sharp instruments. All this, increases the risk of exposure to harmful microorganisms.²



The risk of glove failure rate correlates with the duration of continuous operating time. Overall, the consensus of study results recommends changing gloves no later than 1½ to 2 hours.⁴




During surgical procedures, a greater number of instruments, surgical equipment, technique and reduced surgical experience increases the risk of glove failure rate.³




The exposure to mechanical⁵ and chemical⁶ stress have shown to weaken and degrade the glove overtime. Change gloves if you notice any discoloration, and/or change in shape.⁷

All these factors influence glove integrity and increase perforation risk.

THE RISKS OF SINGLE GLOVING



THE BENEFITS OF DOUBLE GLOVING



Double gloving **reduces risk of exposure** to patient blood by **as much as 87%** when the outer glove is perforated.¹³



Two layers of gloves **reduce the transmission of blood volume** by as much as **95%** thereby reducing the viral load in the event of a contaminated percutaneous injury.¹³



Two layers of gloves can alert the surgeon to glove failure, detecting **77% of perforations versus** only **21%** with single gloves.¹⁵



Wearing a colored under glove increases identification of **perforations by up to 86%**.¹⁴ Colored under gloves also **reduce time to awareness from 67 seconds to 42 seconds**.¹⁸

THE IMPORTANCE OF GLOVE CHANGE



In summary, one major factor for glove failure is wear time. **Glove defect rates are exponentially correlated with longer wear time.**



In a study by Tlilli et al. (2017), it was reported that **significantly higher perforation rates** occurred when the **gloves were worn for over 90 minutes.**¹⁵ AORN, ACORN, AfPP and many other organizations recommend routine glove change. The range averages between **60 - 150 minutes, with 90 minutes being most common.**

WHY SURGICAL STAFF MIGHT NOT PRACTICE DOUBLE GLOVING

Some HCWs claim that dexterity and ability to safely handle and use instruments is compromised or even diminished with the addition of a second layer of gloves. Multiple studies investigating tactility and sensation, both objectively and subjectively, concluded there is **no negative impact on manual dexterity and tactile sensation associated with the use of double gloves.**¹⁶

A range of factors influence and impact PPE-related behaviors. Compliance can be dependent on¹⁷:

Individual Factors

- Knowledge
- Beliefs
- Attitudes
- Experience
- Perception of risk

Environmental Factors

- Availability of glove types/styles
- Cost
- Perceived time

Organizational Factors

- Administrative expectations
- Performance feedback
- Workplace policies



For Healthcare worker and patient safety, always consider the risk when single gloving.
Make double gloving your gold standard for optimal protection.
To learn about the benefits of double gloving, go to www.ansell.com/AnsellCARES.

References:

1. Makama JG, Okeme IM, Makama EJ, Ameh EA. Glove Perforation Rate in Surgery: A Randomized, Controlled Study to Evaluate the Efficacy of Double Gloving. *Surg Infect (Larchmt)*. 2016;17(4):436-442.
2. Jagger J, Berguer R, Phillips EK, Parker G, Gomaa AE. Increase in sharps injuries in surgical settings versus nonsurgical settings after passage of national needlestick legislation. *J Am Coll Surg*. 2010;210(4):496-502.
3. Matsuoka S, Kondo T, Seishima R, Okabayashi K, Tsuruta M, Shigeta K, Ishida T, Hasegawa H, Kitagawa Y. Surgical glove perforation during laparoscopic colorectal procedures. *Surg Endosc*. 2022 May;36(5):3489-3494.
4. Partecke LI, Goerdts A-M, Langner I, et al. Incidence of Micro-perforation for Surgical Gloves Depends on Duration of Wear. *Infection Control & Hospital Epidemiology*. 2009;30(5):409-414
5. Carter AH, Casper DS, Parvizi J, Austin MS. A prospective analysis of glove perforation in primary and revision total hip and total knee arthroplasty. *J Arthroplasty*. 2012 Aug;27(7):1271-5
6. Sabino Ma, Ajami D., Physicochemical, Mechanical, and Biological Properties of Bone Cements Prepared with Functionalized Methacrylates, *Journal of Biomaterial applications*, 19(4):147-61
7. Association of Perioperative of Registered nurses AORN. (2020). Guideline for sterile technique. In: *Guidelines for Perioperative Practice*. Denver, CO: AORN, Inc; 2020.
8. Driever R, Beie M, Schmitz E, et al. Surgical glove perforation in cardiac surgery. *Thorac Cardiovasc Surg* 2001; 49:328-330.
9. Wilburn SQ, Eijkemans G. Preventing needlestick injuries among healthcare workers: a WHO-ICN collaboration. *Int J Occup Environ Health*. 2004;10(4):451-456.
10. International Safety Center. U.S. EPINET Sharps Injury and Blood and Body Fluid Exposure Surveillance Research Group. Sharps Injury Data Report for 2020. Report available at <https://international-safety-center.org/exposure-reports/>. Updated 2020. Accessed February 1, 2022.
11. CDC. Workbook for Designing, Implementing and Evaluating a Sharps Injury Prevention Program. CDC website. https://www.cdc.gov/sharpsafety/pdf/sharpsworkbook_2008.pdf. Published 2008. Accessed January 31, 2022.
12. CDC. Stop Sticks Campaign, Injury report Sharps Injuries: Completing the Injury Report. CDC Website. <https://www.cdc.gov/nora/councils/hcsa/stopsticks/injuryreport.html>. Reviewed February 26, 2019. Accessed February 8, 2022.
13. Berguer R. Key strategies for eliminating sharps injuries during surgery. *AORN Journal*. 2011 Jul;94(1):91-96.
14. Laine T, Aarnio P. How often does glove perforation occur in surgery? Comparison between single gloves and a double-gloving system. *Am J Surg*. 2001;181(6):564-566.
15. Waljee JF, Malay S, Chung KC. Sharps injuries: the risks and relevance to plastic surgeons. *Plast Reconstr Surg*. 2013 Apr;131(4):784-791
16. Fry DE, Harris WE, Kohnke EN, Twomey CL. Influence of double-gloving on manual dexterity and tactile sensation of surgeons. *J Am Coll Surg*. 2010;210(3):325-330.
17. Lipson ME, Deardon R, Switzer NJ, de Gara C, Ball CG, Grondin SC. Practice and attitudes regarding double gloving among staff surgeons and surgical trainees. *Can J Surg*. 2018 Aug;61(4):244-250.
18. Florman S, Burgdorf M, Finigan K, et al., Efficacy of double gloving with intrinsic indicator system, *Surg Infect*;2005; 6(4):385-95.

Supplemental Websites and Additional Information:

1. American Academy of Orthopedic Surgeons (AAOS), <http://www.aaos.org>
2. American College of Surgeons (ACS), <http://www.facs.org>
3. Association for Perioperative Practice (AfPP), <https://www.afpp.org.uk/home>
4. Association of Perioperative Registered Nurses (AORN), <http://www.aorn.org>
5. Association of Surgical Technologists (AST), <http://www.ast.org>
6. Australian College of Perioperative Nurses (ACORN), <https://www.acorn.org.au/>
7. CDC Surgical Site Infection. Guideline for Prevention of Surgical Site Infection. 2017. <https://www.cdc.gov/infectioncontrol/guidelines/ssi/index.html>
8. European Operating Room Nurses Association (EORNA), <https://eorna.eu/>
9. Operating Room Nurses Association of Canada (ORNAC), <https://www.ornac.ca/en/>
10. The World Health Organization Glove Use Information Leaflet 2009. [https://cdn.who.int/media/docs/default-source/integrated-health-services-\(ihs\)/infection-prevention-and-control/hand-hygiene/tools/glove-use-information-leaflet.pdf?sfvrsn=13670aa_10](https://cdn.who.int/media/docs/default-source/integrated-health-services-(ihs)/infection-prevention-and-control/hand-hygiene/tools/glove-use-information-leaflet.pdf?sfvrsn=13670aa_10)

➔ For more information, visit www.ansell.com/AnsellCARES

Ansell, ® and ™ are trademarks owned by Ansell Limited or one of its affiliates. © 2023 All Rights Reserved.

CB.11.23

